

Converse County: Key Surveillance Data



Smoking Rate

Converse County

16%

(11% - 24%)


All Counties

18%

(16% - 20%)

The percentage of adults in Converse County who smoke is not significantly different from all 23 counties combined.


Tried to Quit in the Past Year



The percentage of current and recent smokers³ in Converse County who tried to quit smoking for good in the past year is not significantly different from all 23 counties combined.

Smoked a Whole Cigarette

Before Age 18



The percentage of smokers¹ in Converse County who smoked a whole cigarette for the first time before age 18 is not significantly different from all 23 counties combined.

Want to Quit Using Tobacco for Good

Converse County

63%

(46% - 78%)

All Counties

61%

(56% - 65%)

The percentage of tobacco users⁴ in Converse County who want to quit using tobacco products for good is not significantly different from all 23 counties combined.

Became Regular Smokers

Before Age 18

Converse County

53%

(39% - 66%)

All Counties

45%

(42% - 49%)


The percentage of current and former regular smokers² in Converse County who first smoked at least one cigarette every day for 30 days in a row before age 18 is not significantly different from all 23 counties combined.

1. Includes a) adults who had smoked at least 100 cigarettes in their lifetime and b) smoking experimenters who had ever smoked one whole cigarette.
2. Includes a) current everyday smokers and b) current someday and former smokers who had ever smoked at least one cigarette every day for 30 days in a row.
3. Includes a) current smokers and b) former smokers and smoking experimenters who at least puffed on a cigarette in the past year.
4. Includes a) current smokers, b) former smokers and smoking experimenters who had at least puffed on a cigarette in the past year and had not decided to quit completely and c) current smokeless tobacco users.

Converse County: Key Surveillance Data




Advised to Quit by a Health Professional



The percentage of tobacco users⁵ in Converse County who were advised to quit by a health professional in the past year is not significantly different from all 23 counties combined.

Breathed Secondhand Smoke at Work




The percentage of adults⁶ in Converse County who breathed secondhand smoke at work⁷ in the past week is not significantly different from all 23 counties combined.

5. Includes adults who smoked cigarettes or used smokeless tobacco (including current smokers/users) in the past year *and* had seen a health professional in the past year.

6. Includes adults who are employed or self-employed.


7. Workplace refers to both indoor areas (including the inside of a vehicle) and outdoor areas.

Support a Tax Increase on Cigarettes



The percentage of adults in Converse County who support an increase in the tax on a pack of cigarettes is not significantly different from all 23 counties combined.

Support for Smokefree Indoor Air Laws



The percentages of adults in Converse County who support laws in their community making workplaces, restaurants, and bars smokefree are not significantly different from all 23 counties combined.

Note. Percentages in parentheses and error bars in graphs represent 95% confidence intervals. WYSAC determined statistical significance by non-overlapping confidence intervals of the estimates between a county and all counties combined. For additional information, see the detailed report at <http://wysac.uwyo.edu/wysac/Reports.aspx>.

Wyoming Survey & Analysis Center

University of Wyoming • Dept. 3925
1000 East University Avenue
Laramie, WY 82071
wysac@uwyo.edu • <http://wysac.uwyo.edu>
(307) 766-2189 • Fax: (307) 766-2759

Under contract to
Wyoming Department of Health
Public Health Division
6101 Yellowstone Road, Suite 420
Cheyenne, WY 82002
(307) 777-6340

